


Twenty-second Edition

Harper's Biochemistry



a LANGE medical book

**Robert K. Murray
Daryl K. Granner**

**Peter A. Mayes
Victor W. Rodwell**

a **LANGE** medical book

Harper's Biochemistry

Twenty-second Edition

Robert K. Murray, MD, PhD

*Professor of Biochemistry
University of Toronto*

Daryl K. Granner, MD

*Professor and Chairman
Department of Molecular Physiology and Biophysics
Professor of Medicine
Vanderbilt University
Nashville, Tennessee*

Peter A. Mayes, PhD, DSc

*Reader in Biochemistry
Royal Veterinary College
University of London*

Victor W. Rodwell, PhD

*Professor of Biochemistry
Purdue University
West Lafayette, Indiana*



APPLETON & LANGE

Norwalk, Connecticut / San Mateo, California

1-8385-3640-9

Notice: Our knowledge in clinical sciences is constantly changing. As new information becomes available, changes in treatment and in the use of drugs become necessary. The authors and the publisher of this volume have taken care to make certain that the doses of drugs and schedules of treatment are correct and compatible with the standards generally accepted at the time of publication. The reader is advised to consult carefully the instruction and information material included in the package insert of each drug or therapeutic agent before administration. This advice is especially important when using new or infrequently used drugs.



Copyright © 1990 by Appleton & Lange
A Publishing Division of Prentice Hall
Copyright © 1988 by Appleton & Lange
Previous editions © Lange Medical Publications

All rights reserved. This book, or any parts thereof, may not be used or reproduced in any manner without written permission. For information, address Appleton & Lange, 25 Van Zant Street, East Norwalk, Connecticut 06855.

90 91 92 93 94 / 10 9 8 7 6 5 4 3 2 1

Prentice Hall International (UK) Limited, *London*
Prentice Hall of Australia Pty. Limited, *Sydney*
Prentice Hall Canada, Inc., *Toronto*
Prentice Hall Hispanoamericana, S.A., *Mexico*
Prentice Hall of India Private Limited, *New Delhi*
Prentice Hall of Japan, Inc., *Tokyo*
Simon & Schuster Asia Pte. Ltd., *Singapore*
Editora Prentice Hall do Brasil Ltda., *Rio de Janeiro*
Prentice Hall, *Englewood Cliffs, New Jersey*

ISBN: 0-8385-3640-9
ISSN: 0734-9866

Production Editor: Christine Langan
Designer: Steven Byrum

PRINTED IN THE UNITED STATES OF AMERICA

Preface . .

1. Bloct

2. Biom

3. Wate

Section I

4. Amir

5. Pept

6. Prot

7. Prot

8. Enzy

9. Enzy

10. Enzy

11. Enz

Section

12. Bio

13. Bio


14. Oxi

To locate disulfide bonds, peptides from untreated and from reduced or oxidized protein are separated by 2-dimensional chromatography or by electrophoresis and chromatography (fingerprinting). Visualization with ninhydrin reveals **2 fewer peptides** in the digest from untreated protein and **one new peptide** in the digest from treated protein. With knowledge of the primary structure of these peptides, the positions of disulfide bonds can then be inferred.

PEPTIDES ARE SYNTHESIZED BY AUTOMATED TECHNIQUES


Fig 5-12, which illustrates synthesis of a representative dipeptide A-B by the Merrifield solid-phase technique, summarizes all the reactions required to synthesize a peptide of any desired length. These steps in the procedure are:

1. Block the N-termini of amino acid A (open symbol) and amino acid B (shaded symbol) with the *t*-butoxycarbonyl [*t*-BOC] group (■):



forming *t*-BOC-A and *t*-BOC-B.

2. Activate the carboxyl group of *t*-BOC-B with dicyclohexyl carbodiimide (DCC) (▶):



3. React the carboxyl group of amino acid A (which will become the C-terminal residue of the peptide) with an activated, insoluble polystyrene resin (●).

4. Remove the blocking group from *t*-BOC-A with room temperature trifluoroacetic acid (TFA, F₃C-COOH).

[Note: In practice, steps 3 and 4 may be omitted since resins with any given *t*-BOC-amino acid connected via an ester bond to a phenylacetamidomethyl (PAM) "linker" molecule attached to the polystyrene resin are commercially available.]

5. Condense the activated carboxyl group of *t*-BOC-B with the free amino group of immobilized A.

6. Remove the *t*-BOC blocking group with TFA (see step 4).




Figure 5-11. The overlapping peptide Z is used to deduce that peptides X and Y are present in the original protein in the order X → Y, not Y → X.




Figure 5-12. Symbolic representation of the synthesis of a generic dipeptide by the solid-phase synthesis technique pioneered by Merrifield. See accompanying text for explanations of symbols.

7. Liberate the dipeptide A-B from the resin particle by treating at -2 °C with HF in dichloromethane.

The initial achievements of the Merrifield technique were the synthesis of the A chain (21 residues) and B chain (30 residues) of insulin in 11 days and of the enzyme pancreatic ribonuclease in 18% overall yield. Subsequent improvements have reduced the time for synthesis of a peptide bond to about 1 hour and have increased yields significantly. This has initiated new prospects, not only for confirming *de novo* synthesis of the primary structures of proteins, but for immunology, for producing vaccines and polypeptide hormones, and conceivably also for treating selected in-born errors of metabolism.

REFERENCES

- Allen G: Sequencing of proteins and Peptides. North Holland. 1981.
Bhwon AS: Protein/Peptide Sequence Analysis: Current Methodologies CRC Press, 1988.
Cantor CR, Schimmel PR: Biophysical Chemistry, Part I: The Conformation of Macromolecules. Freeman, 1980.
Dayhoff M (editor): Atlas of Protein Sequence and Structure.

Vol 5: N
ington, J
Doolittle R
Derived
Books.
Hewick RA
liquid so
Chem 1'
Hunkapille
mated n
Hunkapille
226:30
Lipman D
chance